

What should I tell my healthcare provider before starting vigabatrin?

If you or your child has CPS, before taking vigabatrin tell your healthcare provider about all of your medical conditions, including if you or your child:

- have or had an allergic reaction to vigabatrin, such as hives, itching, or trouble breathing.
- have or had any vision problems.
- have or had any kidney problems.
- have or had low red blood cell counts (anemia).
- have or had any nervous or mental illnesses, such as depression, mood problems, thoughts of suicide, or attempts at suicide.
- are breastfeeding or planning to breastfeed. Vigabatrin can pass into breast milk and may harm your baby. Talk to your healthcare provider about the best way to feed your baby if you take vigabatrin.
- are pregnant or plan to become pregnant. Vigabatrin can cause harm to your unborn baby. You and your healthcare provider will have to decide if you should take vigabatrin while you are pregnant.

Pregnancy Registry:

If you become pregnant while taking vigabatrin, talk to your healthcare provider about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334. Information on the registry can also be found at the website <http://www.aedpregnancyregistry.org/>. The purpose of this registry is to collect information about the safety of antiepileptic medicine during pregnancy.

If you are a parent or caregiver whose baby has IS, before giving vigabatrin to your baby, tell your healthcare provider about all of your baby's medical conditions, including if your baby has or ever had:

- an allergic reaction to vigabatrin, such as hives, itching, or trouble breathing.
- any vision problems.
- any kidney problems.

Tell your healthcare provider about all the medicines you or your child take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Vigabatrin and other medicines may affect each other causing side effects.

How should I take vigabatrin?

- Vigabatrin comes as tablets or as packets containing powder for mixing with water to make an oral solution.
- You or your child will receive vigabatrin from a specialty pharmacy.
- Take vigabatrin exactly as your healthcare provider tells you to. Vigabatrin is usually taken 2 times each day.
- Vigabatrin may be taken with or without food.
- Before starting to take vigabatrin, talk to your healthcare provider about what you or your child should do if a vigabatrin dose is missed.
- If you or your child are taking vigabatrin for CPS and the seizures do not improve enough within 3 months, your healthcare provider will stop prescribing vigabatrin.
- If your child is taking vigabatrin for IS and the seizures do not improve within 2 to 4 weeks, your healthcare provider will stop prescribing vigabatrin.
- Do not stop taking vigabatrin suddenly.** This can cause serious problems. Stopping vigabatrin or any seizure medicine suddenly can cause seizures that will not stop (status epilepticus) in people who are being treated for seizures. You should follow your healthcare provider's instructions on how to stop taking vigabatrin.
- Tell your healthcare provider right away about any increase in seizures when vigabatrin treatment is being stopped.** Before your child starts taking vigabatrin, speak to your child's healthcare provider about what to do if your baby misses a dose, vomits, spits up, or only takes part of the dose of vigabatrin.
- Do not stop taking vigabatrin without talking to your healthcare provider.** If vigabatrin improves your (or your child's) seizures, you and your healthcare provider should talk about whether the benefit of taking vigabatrin is more important than the risk of vision loss, and decide if you (or your child) will continue to take vigabatrin.

What should I avoid while taking vigabatrin?

Vigabatrin causes sleepiness and tiredness. Adults taking Vigabatrin should not drive, operate machinery, or perform any hazardous task, unless you and your healthcare provider have decided that you can do these things safely.

What are the possible side effects of vigabatrin?**Vigabatrin can cause serious side effects, including:**

- See** "What is the most important information I should know about vigabatrin?"
- sleepiness and tiredness.** See "What should I avoid while taking vigabatrin?"
- Vigabatrin may cause your baby to be sleepy.** Sleepy babies may have a harder time suckling and feeding, or may be irritable.
- weight gain that happens without swelling.**

The following serious side effects happen in **adults**. It is not known if these side effects also happen in babies who take vigabatrin.

- low red blood cell counts (anemia).**
- nerve problems.** Symptoms of a nerve problem can include numbness and tingling in your toes or feet. It is not known if nerve problems will go away after you stop taking vigabatrin.
- swelling.**

If you or your child has CPS, vigabatrin may make certain types of seizures worse. Tell your healthcare provider right away if your (or your child's) seizures get worse.

The most common side effects of vigabatrin in **adults** include blurred vision, sleepiness, dizziness, problems walking or feeling uncoordinated, shaking (tremor), and tiredness.

The most common side effect of vigabatrin in **children 3 to 16 years of age** is weight gain. Also expect side effects like those seen in adults.

If you are giving vigabatrin to your baby for IS:

Vigabatrin may make certain types of seizures worse. You should tell your baby's healthcare provider right away if your baby's seizures get worse. Tell your baby's healthcare provider if you see any changes in your baby's behavior.

The most common side effects of vigabatrin in **babies** include:

- sleepiness - vigabatrin may cause your baby to be sleepy. Sleepy babies may have a harder time suckling and feeding or may be irritable.
- swelling in the bronchial tubes (bronchitis)
- ear infection
- irritability

Tell your healthcare provider if you or your child have any side effect that bothers you or that does not go away. These are not all the possible side effects of vigabatrin.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800- FDA-1088.

How should I store vigabatrin?

- Store vigabatrin tablets at room temperature, between 68°F to 77°F (20°C to 25°C).

Keep vigabatrin and all medicines out of the reach of children.**General information about the safe and effective use of vigabatrin.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. You can ask your pharmacist or healthcare provider for information about vigabatrin that is written for health professionals. Do not use vigabatrin for a condition for which it was not prescribed. Do not give vigabatrin to other people, even if they have the same symptoms that you have. It may harm them.

What are the ingredients in vigabatrin?

Active Ingredient: vigabatrin, USP

Inactive Ingredients:

- Tablets:** hydroxypropyl methylcellulose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, povidone, polydextrose, sodium starch glycolate, talc, and titanium dioxide.

Distributed by: Carnegie Pharmaceuticals LLC

Delran, NJ 08075, USA

Made In USA

For more information, call 1-732-783-7010.

This Medication Guide has been approved by the U.S. Food and Drug Administration

Revised: 02/2026

Body System Adverse Reaction	All Vigabatrin (N=165)	Placebo (N=104)
Neurologic System Disorders		
Somnolence	6	5
Nystagmus	4	3
Tremor	4	2
Status epilepticus	2	1
Psychiatric Disorders		
Abnormal behavior	7	6
Aggression	6	2
Disorientation	3	0

Safety of vigabatrin for the treatment of refractory CPS in patients 3 years of age is expected to be similar to pediatric patients 3 to 16 years of age.

Adults Spasms

In a randomized, placebo-controlled IS study with a 5-day double-blind treatment phase (n=40), the adverse reactions that occurred in >5% of patients receiving vigabatrin and that occurred more frequently than in placebo patients were somnolence (vigabatrin 45%, placebo 30%), bronchitis (vigabatrin 30%, placebo 15%), ear infection (vigabatrin 10%, placebo 5%), and acute otitis media (vigabatrin 10%, placebo 0%). In a dose response study of low-dose (18-36 mg/kg/day) versus high-dose (100-148 mg/kg/day) vigabatrin, no clear correlation between dose and incidence of adverse reactions was observed. The adverse reactions (>5% in either dose group) are summarized in Table 7.

Table 7. Adverse Reactions in a Placebo-Controlled Trial in Patients with Infantile Spasms

Body System Adverse Reaction	Vigabatrin Low Dose (N=14)	Vigabatrin High Dose (N=10)
Eye Disorders (other than field or acuity changes)		
Strabismus	5	5
Conjunctivitis	5	2
Gastrointestinal Disorders		
Vomiting	14	20
Constipation	14	12
Diarrhea	13	12
General Disorders		
Fever	29	19
Infections		
Upper respiratory tract infection	51	46
Otitis media	44	30
Viral infection	20	19
Pneumonia	13	11
Candidiasis	8	3
Ear infection	7	14
Gastroenteritis viral	6	5
Sinusitis	5	9
Urinary tract infection	5	9
Influenza	5	3
Croup infections	5	1
Metabolism & Nutrition Disorders		
Decreased appetite	9	7
Neurologic System Disorders		
Sedation	19	17
Somnolence	17	19
Status epilepticus	6	4
Lethargy	5	7
Convulsion	4	7
Hypotonia	4	6
Psychiatric Disorders		
Irritability	16	23
Insomnia	10	12
Respiratory Disorders		
Nasal congestion	13	4
Cough	3	8
Skin and Subcutaneous Tissue Disorders		
Rash	8	11

Metabolism & Nutrition Disorders

Decreased appetite

Neurologic System Disorders

Sedation

Somnolence

Status epilepticus

Lethargy

Convulsion

Hypotonia

Psychiatric Disorders

Irritability

Insomnia

Respiratory Disorders

Nasal congestion

Cough

Skin and Subcutaneous Tissue Disorders

Rash

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of vigabatrin. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. Adverse reactions are categorized by system organ class.

Birth Defects: Congenital cardiac defects, congenital external ear anomaly, congenital hemangioma, congenital hypothyroidism, congenital male genital malformation, congenital oral malformation, congenital vesicoureteric reflux, dental/canal anomaly, dysmorphism, fetal anticonvulsant syndrome, hamamans, hip dysplasia, limb malformation, limb reduction defect, low ear ears, renal aplasia, retinitis pigmentosa, supernumerary nipple, talipes

Ear Disorders: Deafness

Endocrine Disorders: Delayed puberty

Gastrointestinal Disorders: Gastrointestinal hemorrhage, esophagitis

General Disorders: Developmental delay, facial edema, malignant hyperthermia, multi-organ failure

Hepatobiliary Disorders: Cholestasis

Nervous System Disorders: Dystonia, encephalopathy, hypertension, hypotonia, muscle spasticity, myoclonus, optic neuritis, dyskinesia

Respiratory Disorders: Acute psychosis, apathy, delirium, hypomania, neonatal apnea, psychotic disorder, respiratory disorders, Laryngeal edema, pulmonary embolism, respiratory failure, stridor

Skin and Subcutaneous Tissue Disorders: Angioedema, maculo-papular rash, pruritus, Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), alopecia

7 DRUG INTERACTIONS

7.1 Antiepileptic Drugs

Phenytoin

Although phenytoin dose adjustments are not routinely required, dose adjustment of phenytoin should be considered if clinically indicated, since vigabatrin may cause a moderate reduction in total phenytoin plasma levels (see Clinical Pharmacology (12.3)).

Clonazepam

Vigabatrin may moderately increase the C_{ss} of clonazepam resulting in an increase of clonazepam-associated adverse reactions (see Clinical Pharmacology (12.3)).

Other AEDs

There are no clinically significant pharmacokinetic interactions between vigabatrin and either phenobarbital or sodium valproate. Based on population pharmacokinetics, carbamazepine, clobazepam, primidone, and sodium valproate appear to have no effect on plasma concentrations of vigabatrin (see Clinical Pharmacology (12.3)).

7.2 Oral Contraceptives

Vigabatrin is unlikely to affect the efficacy of steroid oral contraceptives (see Clinical Pharmacology (12.3)).

7.3 Drug-Laboratory Test Interactions

Vigabatrin decreases alanine transaminase (ALT) and aspartate transaminase (AST) plasma activity in up to 80% of patients. In some patients, these enzymes become undetectable. The suppression of ALT and AST activity by vigabatrin may preclude the use of these markers, especially ALT, to detect early hepatic injury.

Vigabatrin may increase the amount of amino acids in the urine, possibly leading to a false positive test for certain rare genetic metabolic diseases (e.g., alpha aminoaciduria).

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to AEDs, including vigabatrin, during pregnancy. Encourage women who are taking vigabatrin during pregnancy to enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry. This can be done by calling the toll-free number 1-888-233-2334 or visiting the website, <http://www.aedpregnancyregistry.org/>. This must be done by the patient herself.

Risk Summary

There are no adequate data on the developmental risk associated with the use of vigabatrin in pregnant women. Limited available data from case reports and cohort studies pertaining to vigabatrin use in pregnant women have not established a drug-associated risk of major birth defects, miscarriage, or adverse maternal

or fetal outcomes. However, based on animal data, vigabatrin use in pregnant women may result in fetal harm.

When administered to pregnant animals, vigabatrin produced developmental toxicity, including an increase in fetal malformations and offspring neurobehavioral and neurohistopathological effects. At clinically relevant doses, developmental neurotoxicity was observed in rats treated with vigabatrin during a period of postnatal development corresponding to the third trimester of human pregnancy (see Data).

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2%-4% and 15%-20%, respectively. The background risk of major birth defects and miscarriage for the indicated population is unknown.

Data**Animal Data**

Administration of vigabatrin (oral doses of 50 to 200 mg/kg/day) to pregnant rabbits throughout the period of organogenesis was associated with an increased incidence of malformations (cleft palate) and embryofetal death. These findings were observed in two separate studies. The no-effect dose for adverse effects on embryofetal development in rabbits (100 mg/kg/day) is approximately 1/2 the maximum recommended human dose (MRHD) of 3 g/day on a body surface area (mg/m²) basis. In rats, oral administration of vigabatrin (50, 100, or 150 mg/kg/day) throughout organogenesis resulted in decreased fetal body weights and increased incidences of fetal anatomic variations. The no-effect dose for adverse effects on embryo-fetal development in rats (50 mg/kg/day) is approximately 1/5 the MRHD on a mg/m² basis. Oral administration of vigabatrin (50, 100, 150 mg/kg/day) to rats from the latter part of pregnancy through weaning produced long-term neurohistopathological (hippocampal vacuolation) and neurobehavioral (consolidation) abnormalities in the offspring. A no-effect dose for developmental neurotoxicity in rats was not established; the low effect dose (50 mg/kg/day) is approximately 1/5 the MRHD on a mg/m² basis.

In a published study, vigabatrin (300 to 450 mg/kg) was administered by intraperitoneal injection to a mutant mouse strain on a single day during organogenesis (day 8, 9, 10, 11, or 12). An increase in fetal malformations (including cleft palate) was observed at both doses.

Oral administration of vigabatrin (5, 15, or 50 mg/kg/day) to young rats during the neonatal and juvenile periods of development (postnatal days 4-63) produced neurobehavioral (convulsions, neuroleptic impairment, learning deficits) and neurohistopathological (brain vacuolation, decreased myelination, and retinal dysplasia) abnormalities in treated animals. The early postnatal period in rats is generally thought to correspond to late pregnancy in humans in terms of brain development. The no-effect dose for developmental neurotoxicity in juvenile rats (5 mg/kg/day) was associated with plasma vigabatrin exposures (AUC) less than 1/20 of those measured in pediatric patients receiving an oral dose of 50 mg/kg.

8.2 Lactation**Risk Summary**

Vigabatrin is excreted in human milk. The effects of vigabatrin on the breastfed infant and on milk production are unknown. Because of the potential for serious adverse reactions from vigabatrin in nursing infants, breastfeeding is not recommended. If a breastfeeding infant is taking vigabatrin, observe for any potential adverse effects (see Warnings and Precautions (5.1, 5.3, 5.4, 5.8)).

8.4 Pediatric Use

The safety and effectiveness of vigabatrin as adjunctive treatment of refractory complex partial seizures in pediatric patients 2 to 16 years of age have been established and is supported by three double-blind, placebo-controlled studies in patients 3 to 16 years of age, adequate and well-controlled studies in adult patients, pharmacokinetic data from patients 2 years of age and older, and additional safety information (3 years to 16 years of age). US Patients for children and adolescents (10 to 16 years of age), and 10-15 hours for adults. Following administration in this population varies according to age group and is weight-based (see Dosage and Administration (2.2)). Adverse reactions in this pediatric population are similar to those observed in the adult population (see Adverse Reactions (6.1)). The safety and effectiveness of vigabatrin as monotherapy for pediatric patients with infantile spasms (1 month to 2 years of age) have been established (see Dosage and Administration (2.3) and Clinical Studies (14.2)).

Safety and effectiveness as adjunctive treatment of refractory complex partial seizures in pediatric patients below the age of 2 and as monotherapy for the treatment of infantile spasms in pediatric patients below the age of 1 month have not been established.

Duration of therapy for infantile spasms was evaluated in a post hoc analysis of a Canadian Pediatric Epilepsy Network (CPEN) study of developmental outcomes in infantile spasms patients. This analysis suggests that a total duration of 6 months of vigabatrin therapy is adequate for the treatment of infantile spasms. However, prescribers must use their clinical judgment as to the most appropriate duration of use (see Clinical Studies (14.2)).

Abnormal MRI signal changes and Intracranial Edema (IME) in infants and young children being treated with vigabatrin have been observed (see Warnings and Precautions (5.3, 5.4)).

Juvenile Animal Toxicity Data

Oral administration of vigabatrin (5, 15, or 50 mg/kg/day) to young rats during the neonatal and juvenile periods of development (postnatal days 4-63) produced neurobehavioral (convulsions, neuroleptic impairment, learning deficits) and neurohistopathological (brain gray matter vacuolation, decreased myelination, and retinal dysplasia) abnormalities. The no-effect dose for developmental neurotoxicity in juvenile rats (the lowest dose tested) was associated with plasma vigabatrin exposures (AUC) substantially less than those measured in pediatric patients at recommended doses. In dogs, oral administration of vigabatrin (30 or 100 mg/kg/day) during selected periods of juvenile development (postnatal days 22-112) produced neurohistopathological (brain gray matter vacuolation) and neurobehavioral effects of vigabatrin were not assessed in the juvenile dog. A no-effect dose for neurohistopathology was not established in juvenile dogs; the lowest effect dose (30 mg/kg/day) was associated with plasma vigabatrin exposures lower than those measured in pediatric patients at recommended doses (see Warnings and Precautions (5.4)).

8.5 Geriatric Use

Clinical studies of vigabatrin did not include sufficient numbers of patients aged 65 and over to determine whether they responded differently from younger patients.

Vigabatrin is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

Oral administration of a single dose of 1.5 g of vigabatrin to elderly (>65 years) patients with reduced creatinine clearance (<50 mL/min) was associated with moderate to severe sedation and hypotension in 4 of 5 patients, lasting up to 5 days. The renal clearance of vigabatrin was 36% lower in healthy elderly subjects (>65 years) than in young healthy males. Adjustment of dose or frequency of administration should be considered. Such patients may respond to a lower maintenance dose (see Dosage and Administration (2.4) and Clinical Pharmacology (12.3)).

Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

8.6 Renal Impairment

Dose adjustment, including starting at a lower dose, is necessary in pediatric patients 2 years of age and older and adults with mild (creatinine clearance >30 to 50 mL/min), moderate (creatinine clearance >30 to 50 mL/min) and severe (creatinine clearance >10 to 30 mL/min) renal impairment (see Dosage and Administration (2.4) and Clinical Pharmacology (12.3)).

9 DRUG ABUSE AND DEPENDENCE**9.1 Controlled Substance**

Vigabatrin is not a controlled substance.

9.2 Abuse

Vigabatrin did not produce adverse events or overt behaviors associated with abuse when administered to humans or animals. It is not possible to predict the extent to which a CNS active drug will be misused, diverted, and/or abused once marketed. Consequently, physicians should carefully evaluate patients for history of drug abuse and follow such patients closely, observing them for signs of misuse or abuse of vigabatrin (e.g., incrementation of dose, drug-seeking behavior).

9.3 Dependence

Following chronic administration of vigabatrin to animals, there were no apparent withdrawal signs upon drug discontinuation. However, as with all AEDs, vigabatrin should be withdrawn gradually to minimize increased seizure frequency (see Warnings and Precautions (5.6)).

10 OVERDOSAGE

10.1 Signs, Symptoms, and Laboratory Findings of Overdose

Confusion and/or suspected vigabatrin overdoses have been reported during clinical trials and in post marketing surveillance. No vigabatrin overdoses resulted in death. When reported, the vigabatrin dose ingested ranged from 3 g to 80 g, but most were between 7.5 g and 30 g. Nearly half the cases involved multiple drug ingestions including carbamazepine, barbiturates, benzodiazepines, lamotrigine, valproic acid, acetaminophen, and/or chlorpheniramine.

Coma, unconsciousness, and/or drowsiness were described in the majority of cases of vigabatrin overdose. Other less commonly reported symptoms included vertigo, psychosis, apnea or respiratory depression, bradycardia, agitation, irritability, confusion, headache, hypotension, abnormal behavior, increased seizure activity, status epilepticus, and speech disorder. These symptoms resolved with supportive care.

10.2 Management of Overdose

There is no specific antidote for vigabatrin overdose. Standard measures to remove unabsorbed drug should be used, including emesis or gastric lavage. Supportive measures should be employed, including monitoring of vital signs and observation of the clinical status of the patient.

In an *in vitro* study, activated charcoal did not significantly adsorb vigabatrin.

Vigabatrin was negative in *in vitro* Ames, CHO/clastogenicity, mammalian cell forward gene mutation, chromosomal aberration in rat lymphocytes and in *in vivo* mouse bone marrow micronucleus assays.

No adverse effects on male or female fertility were observed in rats at oral doses up to 150 mg/kg/day (approximately 1/2 the MRHD of 3 g/day on a mg/m² basis for refractory complex partial seizures).

11 DESCRIPTION

Vigabatrin, USP is an oral antiepileptic drug and is available as white film-coated 500 mg tablets.

The chemical name, a racemate consisting of two enantiomers, is (4S)-4-amino-5-hexenoic acid. The molecular formula is C₆H₁₁NO₂, and the molecular weight is 129.16. It has the following structural formula:



Vigabatrin, USP is a white or almost white powder which is freely soluble in water, slightly soluble in methyl alcohol, very slightly soluble in ethyl alcohol and chloroform, and insoluble in toluene and hexane. The pH of a 1% aqueous solution is about 6.5. The n-octanol/water partition coefficient of vigabatrin is about 0.071 (log P = -1.84) at physiological pH. Vigabatrin, USP melts with decomposition in a 5-degree range within the temperature interval of 171°C to 176°C. The dissociation constants (pK_a) of vigabatrin are 4 and 9.7 at room temperature (25°C).

Each vigabatrin tablet, USP contains 500 mg of vigabatrin, USP. The inactive ingredients are hydroxypropyl methylcellulose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, povidone, polydextrose, sodium starch glycolate, talc, and titanium dioxide.

FDA approved dissolution test specifications other from USP.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The precise mechanism of vigabatrin's anti-seizure effect is unknown, but it is believed to be the result of its action as an irreversible inhibitor of γ-aminobutyric acid transaminase (GABA-T). The enzyme responsible for the metabolism of the inhibitory neurotransmitter GABA. This action results in increased levels of GABA in the central nervous system.

No direct correlation between plasma concentration and efficacy has been established. The duration of drug effect is presumed to be dependent on the rate of enzyme re-synthesis rather than on the rate of elimination of the drug from the systemic circulation.

12.2 Pharmacodynamics

Effects on Electroencephalogram

There is no indication of a QT/